References

1 Berkow R, ed. *The Merck Manual of Medical Information.* Whitehouse Station, NJ: Merck Research Laboratories, 1997.

2 *American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition*, *Text Revision*. Washington, DC: American Psychiatric Association, 2000.

3 Béland Y, Dufour J, Gravel R. Sample design of the Canadian Mental Health Survey. *Proceedings of the Survey Methods Section, 2001.* Vancouver: Statistical Society of Canada, 2001: 93-8.

4 Statistics Canada. *Canadian Community Health Survey (CCHS) - Mental Health and Well-being - Cycle 1.2.* Available at: http://www.statcan.ca/english/concepts/health/cycle1-2/index. htm. Accessed May 11, 2004.

5 Yonkers KA, Zlotnick C, Allsworth J, et al. Is the course of panic disorder the same in women and men? *American Journal of Psychiatry* 1998; 155(5): 596-602.

6 Wittchen EU, Essau CA. Epidemiology of panic disorder: progress and unresolved issues. *Journal of Psychiatric Research* 1993; 27 (Suppl 1): 47-68.

7 Rouillon F. Epidemiology of panic disorder. *Human Psychopharmacology* 1997; 12: S7-S12.

8 Ballenger JC, Lydiard RB. Panic disorder: results of a patient survey. *Human Psychopharmacology* 2004; 12: S27-S33.

9 Béland Y. Canadian Community Health Survey—Methodological overview. *Health Reports* (Statistics Canada, Catalogue 82-003) 2002; 13(3): 9-14.

10 Rao JNK, Wu CFJ, Yue K. Some recent work on resampling methods for complex surveys. *Survey Methodology* (Statistics Canada, Catalogue 12-001) 1992; 18(2): 209-17.

11 Rust KF, Rao JNK. Variance estimation for complex surveys using replication techniques. *Statistical Methods in Medical Research* 1996; 5: 281-310.

12 Alonso J, Angermeyer MC, Bernet S, et al. 12-month comorbidity patterns and associated factors in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. *Acta Psychiatrica Scandinavica Supplementum* 2004; 420: 28-37.

13 Goisman RM, Warshaw MG, Peterson LG, et al. Panic, agoraphobia, and panic disorder with agoraphobia. Data from a multicenter anxiety disorders study. *The Journal of Nervous and Mental Disease* 1994; 182(2): 72-9.

14 Andersch S, Hetta J. A 15-year follow-up study of patients with panic disorder. *European Psychiatry: The Journal of the European College of Neuropsychopharmacology* 2003; 18(8): 401-8.

15 Starcevic V, Uhlenhuth EH, Kellner R, et al. Patterns of comorbidity in panic disorder and agoraphobia. *Psychiatry Research* 1992; 42(2): 171-83.

16 Eaton WW, Kessler RC, Wittchen HU, et al. Panic and panic disorder in the United States. *American Journal of Psychiatry* 1994; 151(3): 413-20.

17 Alonso J, Angermeyer MC, Bernert S, et al. Prevalence of mental disorders in Europe: results from the European Study of Epidemiology of Mental Disorders (ESEMED) project. *Acta Psychiatrica Scandinavica Supplementum* 2004 (420): 21-7.

18 Carlbring P, Gustafsson H, Ekselius L, et al. 12-month prevalence of panic disorder with or without agoraphobia in the Swedish general population. *Social Psychiatry and Psychiatric Epidemiology* 2002; 37(5): 207-11.

19 Weissman MM, Bland RC, Canino GJ, et al. The cross-national epidemiology of panic disorder. *Archives of General Psychiatry* 1997; 54(4): 305-9.

20 Grant B, Katon W, Beitman B. Panic disorder. *Journal of Family Practice* 1983; 17(5): 907-14.

21 Marcaurelle R, Bélanger C, Marchand A. Marital relationship and the treatment of panic disorder with agoraphobia: a critical review. *Clinical Psychology Review* 2003; 23(2): 247-76.

22 Keller MB, Yonkers KA, Warshaw MG, et al. Remission and relapse in subjects with panic disorder and panic with agoraphobia: a prospective short-interval naturalistic follow-up. *The Journal of Nervous and Mental Disease* 1994; 182(5): 290-6.

23 Kessler RC, Stang PE, Wittchen HU et al. Lifetime panic-depression comorbidity in the National Comorbidity Survey of the United States. *Archives of General Psychiatry* 1998; 55(9): 801-8.

24 Iketani T, Kiriike N, Stein MB, et al. Personality disorder comorbidity in panic disorder patients with or without current major depression. *Depression and Anxiety* 2002; 15(4): 176-82.

25 Hino T, Takeuchi T, Yamanouchi N. A 1-year follow-up study of coping in patients with panic disorder *Comprehensive Psychiatry* 2002; 43(4): 279-84.

26 Yamada K, Fujii I, Akiyoshi J, et al. Coping behaviour in patients with panic disorder. *Psychiatry Clinical Neuroscience* 2004; 58(2): 173-8.

27 Vitaliano PP, Katon W, Russo J, et al. Coping as an index of illness behaviour in panic disorder. *Journal of Nervous and Mental Disease* 1987; 175(2): 78-84.

28 Turgeon L, Marchand A, Dupuis G. Clinical features in panic disorder with agoraphobia: a comparison of men and women. *Journal of Anxiety Disorders* 1998; 12(6): 539-53.

29 Starcevic V, Djordjevic A, Latas M, et al. Characteristics of agoraphobia in women and men with panic disorder with agoraphobia. *Depression and Anxiety* 1998; 8(1): 8-13.

30 Sachs GS, Amering P, Berger P, et al. Gender-related disabilities in panic disorder. *Archives of Women's Mental Health* 2002; 4: 121-7.

31 Dannon PN, Lowengrub K, Amaiz R, et al. Comorbid cannabis use and panic disorder : short-term and long-term follow-up. *Human Psychopharmacology: Clinical & Experimental* 2004; 19(2): 97.

32 Katon WJ, Von Korff M, Lin E. Panic disorder: relationship to high medical utilization, unexplained physical symptoms and medical costs. *The American Journal of Medicine.* 1992; 92(1A): 7S-11S.

33 Zaubler TS, Katon W. Panic disorder and medical comorbidity: a review of the medical and psychiatric literature. *Bulletin of the Menninger Clinic* 1996; 60(2 Suppl A): A12-A38.

34 Katerndahl DA, Realini JP. Use of health care services by persons with panic symptoms. *Psychiatric Services* 1997; 48(8): 1027-32.

35 Katerndahl DA. Predictors and outcomes in people told they have panic attacks. *Depression and Anxiety* 2003; 17(2): 98-100.

36 Katon W. Panic disorder: epidemiology, diagnosis, and treatment in primary care. *Journal of Clinical Psychiatry* 1986; 47 (Suppl): 21-30.

37 Yingling KW, Wulsin LR, Arnold LM, et al. Estimated prevalence of panic disorder and depression among consecutive patients seen in an emergency department with acute chest pain. *Journal of General Internal Medicine* 1993; 8(5): 231-5.

38 Katon W. Panic disorder: relationship to high medical utilization, unexplained physical symptoms, and medical costs. *Journal of Clinical Psychiatry* 1996; 57 (Suppl 10): 11-8.

39 Olfson M, Kessler RC, Berglund PA, et al. Psychiatric disorder onset and first treatment contact in the United States and Ontario. *American Journal of Psychiatry* 1998; 155(10): 1415-22.

References

1 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision*. Washington, DC: American Psychiatric Association, 2000.

2 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Third Edition, Text Revision*. Washington DC: American Psychiatric Association, 1987.

3 Walters EE, Kessler RC, Nelson CB, et al. Scoring the World Health Organization's Composite International Diagnostic Interview Short Form (CIDI-SF). www.who.int/msa/cidi/CIDISFScoringMemo12-03-02.pdf.